Appendix E: Metals Trending Graphs Part 1
Alberta/Saskatchewan Border



Figure E1 Battle River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.26
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 11.26
Adjusted Kruskal-Wallis statistic (H') = 11.26



Figure E2 Battle River: Aluminum Dissolved



Figure E3 Battle River: Aluminum Dissolved



Figure E4 Battle River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITY at the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.452
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E5 Battle River: Aluminum Total

Sen's Slope Estimator



Figure E6 Battle River: Aluminum Total



Figure E7 Battle River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 43.92
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 43.92
Adjusted Kruskal-Wallis statistic (H') = 43.92



Figure E8 Battle River: Arsenic Dissolved



Figure E9 Battle River: Arsenic Dissolved



Figure E10 Battle River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.8
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E11 Battle River: Arsenic Total

Seasonal Kendall



Figure E12 Battle River: Arsenic Total



Figure E13 Battle River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 25.96
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E14 Battle River: Barium Dissolved



Figure E15 Battle River: Barium Dissolved



Figure E16 Battle River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 33.21
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E17 Battle River: Barium Total

Seasonal Kendall



Figure E18 Battle River: Barium Total



Figure E19 Battle River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4279
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.4279
Adjusted Kruskal-Wallis statistic (H') = 0.4279



Figure E20 Battle River: Beryllium Dissolved



Figure E21 Battle River: Beryllium Dissolved



Figure E22 Battle River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.05818
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.05818
Adjusted Kruskal-Wallis statistic (H') = 0.05818



Figure E23 Battle River: Beryllium Total

Sen's Slope Estimator



Figure E24 Battle River: Beryllium Total



Figure E25 Battle River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.41
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E26 Battle River: Boron Dissolved



Figure E27 Battle River: Boron Dissolved



Figure E28 Battle River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.918
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E29 Battle River: Boron Total

Seasonal Kendall



Figure E30 Battle River: Boron Total



Figure E31 Battle River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.996
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E32 Battle River: Cadmium Dissolved



Figure E33 Battle River: Cadmium Dissolved



Figure E34 Battle River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.511
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E35 Battle River: Cadmium Total

Seasonal Kendall



Figure E36 Battle River: Cadmium Total



Figure E37 Battle River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.0001189
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.0001189
Adjusted Kruskal-Wallis statistic (H') = 0.0001189



Figure E38 Battle River: Chromium Dissolved



Figure E39 Battle River: Chromium Dissolved



Figure E40 Battle River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.002147
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.002147
Adjusted Kruskal-Wallis statistic (H') = 0.002147



Figure E41 Battle River: Chromium Total

Sen's Slope Estimator



Figure E42 Battle River: Chromium Total



Figure E43 Battle River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1866
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E44 Battle River: Cobalt Dissolved



Figure E45 Battle River: Cobalt Dissolved



Figure E46 Battle River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2556
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E47 Battle River: Cobalt Total

Sen's Slope Estimator



Figure E48 Battle River: Cobalt Total



Figure E49 Battle River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.94
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 8.94
Adjusted Kruskal-Wallis statistic (H') = 8.94



Figure E50 Battle River: Copper Dissolved



Figure E51 Battle River: Copper Dissolved



Figure E52 Battle River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.512
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.512
Adjusted Kruskal-Wallis statistic (H') = 2.512



Figure E53 Battle River: Copper Total

Sen's Slope Estimator



Figure E54 Battle River: Copper Total



Figure E55 Battle River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.959
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E56 Battle River: Iron Dissolved



Figure E57 Battle River: Iron Dissolved



Figure E58 Battle River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.038
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E59 Battle River: Iron Total

Seasonal Kendall



Figure E60 Battle River: Iron Total



Figure E61 Battle River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.178
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E62 Battle River: Lead Dissolved



Figure E63 Battle River: Lead Dissolved



Figure E64 Battle River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1384
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E65 Battle River: Lead Total

Sen's Slope Estimator



Figure E66 Battle River: Lead Total



Figure E67 Battle River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.216
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E68 Battle River: Lithium Dissolved



Figure E69 Battle River: Lithium Dissolved



Figure E70 Battle River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.37
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E71 Battle River: Lithium Total



Figure E72 Battle River: Lithium Total

Time Series



Figure E73 Battle River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 43.27
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E74 Battle River: Manganese Dissolved



Figure E75 Battle River: Manganese Dissolved



Figure E76 Battle River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 39.54
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E77 Battle River: Manganese Total

Seasonal Kendall



Figure E78 Battle River: Manganese Total



Figure E79 Battle River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8362
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E80 Battle River: Molybdenum Dissolved



Figure E81 Battle River: Molybdenum Dissolved



Figure E82 Battle River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.651
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.651
Adjusted Kruskal-Wallis statistic (H') = 1.651



Figure E83 Battle River: Molybdenum Total

Sen's Slope Estimator



Figure E84 Battle River: Molybdenum Total



Figure E85 Battle River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.3292
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E86 Battle River: Nickel Dissolved



Figure E87 Battle River: Nickel Dissolved



Figure E88 Battle River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.953
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E89 Battle River: Nickel Total

Seasonal Kendall



Figure E90 Battle River: Nickel Total



Figure E91 Battle River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.831
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 5.831
Adjusted Kruskal-Wallis statistic (H') = 5.831



Figure E92 Battle River: Selenium Dissolved



Figure E93 Battle River: Selenium Dissolved



Figure E94 Battle River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITY at the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.3354
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.3354
Adjusted Kruskal-Wallis statistic (H') = 0.3354



Figure E95 Battle River: Selenium Total

Sen's Slope Estimator



Figure E96 Battle River: Selenium Total

Time Series



Figure E97 Battle River: Silver Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.842
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 5 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.842
Adjusted Kruskal-Wallis statistic (H') = 2.842



Figure E98 Battle River: Silver Dissolved



Figure E99 Battle River: Silver Dissolved



Figure E100 Battle River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.107
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 8 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.107
Adjusted Kruskal-Wallis statistic (H') = 0.107



Figure E101 Battle River: Silver Total

Sen's Slope Estimator



Figure E102 Battle River: Silver Total



Figure E103 Battle River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5138
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.5138
Adjusted Kruskal-Wallis statistic (H') = 0.5138



Figure E104 Battle River: Thallium Dissolved



Figure E105 Battle River: Thallium Dissolved



Figure E106 Battle River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1384
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.1384
Adjusted Kruskal-Wallis statistic (H') = 0.1384



Figure E107 Battle River: Thallium Total

Sen's Slope Estimator



Figure E108 Battle River: Thallium Total



Figure E109 Battle River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.98
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E110 Battle River: Uranium Dissolved



Figure E111 Battle River: Uranium Dissolved



Figure E112 Battle River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.11
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E113 Battle River: Uranium Total

Seasonal Kendall



Figure E114 Battle River: Uranium Total



Figure E115 Battle River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 56.2
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E116 Battle River: Vanadium Dissolved



Figure E117 Battle River: Vanadium Dissolved



Figure E118 Battle River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.08
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E119 Battle River: Vanadium Total

Seasonal Kendall



Figure E120 Battle River: Vanadium Total



Figure E121 Battle River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.56
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E122 Battle River: Zinc Dissolved



Figure E123 Battle River: Zinc Dissolved



Figure E124 Battle River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.35
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.35
Adjusted Kruskal-Wallis statistic (H') = 3.35



Figure E125 Battle River: Zinc Total

Sen's Slope Estimator



Figure E126 Battle River: Zinc Total



Figure E127 Beaver River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.814
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E128 Beaver River: Aluminum Dissolved



Figure E129 Beaver River: Aluminum Dissolved



Figure E130 Beaver River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.567
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.567
Adjusted Kruskal-Wallis statistic (H') = 0.567



Figure E131 Beaver River: Aluminum Total

Sen's Slope Estimator



Figure E132 Beaver River: Aluminum Total



Figure E133 Beaver River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.7
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 18.7
Adjusted Kruskal-Wallis statistic (H') = 18.7



Figure E134 Beaver River: Arsenic Dissolved



Figure E135 Beaver River: Arsenic Dissolved



Figure E136 Beaver River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.725
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E137 Beaver River: Arsenic Total

Seasonal Kendall



Figure E138 Beaver River: Arsenic Total



Figure E139 Beaver River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.68
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E140 Beaver River: Barium Dissolved



Figure E141 Beaver River: Barium Dissolved



Figure E142 Beaver River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 12.54
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E143 Beaver River: Barium Total

Seasonal Kendall



Figure E144 Beaver River: Barium Total



Figure E145 Beaver River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 5 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 7.31
Adjusted Kruskal-Wallis statistic (H') = 7.31



Figure E146 Beaver River: Beryllium Dissolved



Figure E147 Beaver River: Beryllium Dissolved



Figure E148 Beaver River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1484
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 4 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.1484
Adjusted Kruskal-Wallis statistic (H') = 0.1484



Figure E149 Beaver River: Beryllium Total

Sen's Slope Estimator



Figure E150 Beaver River: Beryllium Total



Figure E151 Beaver River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.767
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E152 Beaver River: Boron Dissolved



Figure E153 Beaver River: Boron Dissolved



Figure E154 Beaver River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.011
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E155 Beaver River: Boron Total

Seasonal Kendall



Figure E156 Beaver River: Boron Total



Figure E157 Beaver River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6071
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E158 Beaver River: Cadmium Dissolved



Figure E159 Beaver River: Cadmium Dissolved



Figure E160 Beaver River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.201
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E161 Beaver River: Cadmium Total

Seasonal Kendall



Figure E162 Beaver River: Cadmium Total



Figure E163 Beaver River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.44
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E164 Beaver River: Chromium Dissolved



Figure E165 Beaver River: Chromium Dissolved



Figure E166 Beaver River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.008245
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E167 Beaver River: Chromium Total

Sen's Slope Estimator



Figure E168 Beaver River: Chromium Total



Figure E169 Beaver River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.387
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E170 Beaver River: Cobalt Dissolved



Figure E171 Beaver River: Cobalt Dissolved



Figure E172 Beaver River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.442
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E173 Beaver River: Cobalt Total

Sen's Slope Estimator



Figure E174 Beaver River: Cobalt Total



Figure E175 Beaver River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.471
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E176 Beaver River: Copper Dissolved



Figure E177 Beaver River: Copper Dissolved



Figure E178 Beaver River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2088
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E179 Beaver River: Copper Total

Sen's Slope Estimator



Figure E180 Beaver River: Copper Total



Figure E181 Beaver River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.614
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E182 Beaver River: Iron Dissolved



Figure E183 Beaver River: Iron Dissolved



Figure E184 Beaver River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.399
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E185 Beaver River: Iron Total

Seasonal Kendall



Figure E186 Beaver River: Iron Total



Figure E187 Beaver River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.149
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E188 Beaver River: Lead Dissolved



Figure E189 Beaver River: Lead Dissolved



Figure E190 Beaver River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.3966
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E191 Beaver River: Lead Total

Sen's Slope Estimator



Figure E192 Beaver River: Lead Total

Time Series



Figure E193 Beaver River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.43
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E194 Beaver River: Lithium Dissolved



Figure E195 Beaver River: Lithium Dissolved



Figure E196 Beaver River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 16.84
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E197 Beaver River: Lithium Total

Seasonal Kendall



Figure E198 Beaver River: Lithium Total



Figure E199 Beaver River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 16.43
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E200 Beaver River: Manganese Dissolved



Figure E201 Beaver River: Manganese Dissolved



Figure E202 Beaver River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.08752
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E203 Beaver River: Manganese Total

Sen's Slope Estimator



Figure E204 Beaver River: Manganese Total



Figure E205 Beaver River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.0232
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E206 Beaver River: Molybdenum Dissolved



Figure E207 Beaver River: Molybdenum Dissolved



Figure E208 Beaver River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1135
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E209 Beaver River: Molybdenum Total

Sen's Slope Estimator



Figure E210 Beaver River: Molybdenum Total



Figure E211 Beaver River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.008786
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E212 Battle River: Nickel Dissolved



Figure E213 Beaver River: Nickel Dissolved



Figure E214 Beaver River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.00248
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E215 Beaver River: Nickel Total

Sen's Slope Estimator



Figure E216 Beaver River: Nickel Total



Figure E217 Beaver River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5514
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.5514
Adjusted Kruskal-Wallis statistic (H') = 0.5514



Figure E218 Beaver River: Selenium Dissolved



Figure E219 Beaver River: Selenium Dissolved



Figure E220 Beaver River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.289
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.289
Adjusted Kruskal-Wallis statistic (H') = 4.289



Figure E221 Beaver River: Selenium Total

Seasonal Kendall



Figure E222 Beaver River: Selenium Total



Figure E223 Beaver River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8379
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8379
Adjusted Kruskal-Wallis statistic (H') = 0.8379



Figure E224 Beaver River: Silver Total



Figure E225 Beaver River: Silver Total



Figure E226 Beaver River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8008
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8008
Adjusted Kruskal-Wallis statistic (H') = 0.8008



Figure E227 Beaver River: Thallium Dissolved

Sen's Slope Estimator



Figure E228 Beaver River: Thallium Dissolved



Figure E229 Beaver River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01305
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.01305
Adjusted Kruskal-Wallis statistic (H') = 0.01305



Figure E230 Beaver River: Thallium Total



Figure E231 Beaver River: Thallium Total



Figure E232 Beaver River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.401
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E233 Beaver River: Uranium Dissolved

Sen's Slope Estimator



Figure E234 Beaver River: Uranium Dissolved



Figure E235 Beaver River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.876
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E236 Beaver River: Uranium Total



Figure E237 Beaver River: Uranium Total



Figure E238 Beaver River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.37
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E239 Beaver River: Vanadium Dissolved

Seasonal Kendall



Figure E240 Beaver River: Vanadium Dissolved



Figure E241 Beaver River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.156
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E242 Beaver River: Vanadium Total



Figure E243 Beaver River: Vanadium Total



Figure E244 Beaver River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.363
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.363
Adjusted Kruskal-Wallis statistic (H') = 1.363



Figure E245 Beaver River: Zinc Dissolved

Sen's Slope Estimator



Figure E246 Beaver River: Zinc Dissolved



Figure E247 Beaver River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.539
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E248 Beaver River: Zinc Total



Figure E249 Beaver River: Zinc Total



Figure E250 Cold River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.835
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.835
Adjusted Kruskal-Wallis statistic (H') = 4.835



Figure E251 Cold River: Aluminum Dissolved



Figure E252 Cold River: Aluminum Dissolved



Figure E253 Cold River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.397
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.397
Adjusted Kruskal-Wallis statistic (H') = 1.397



Figure E254 Cold River: Aluminum Total

Sen's Slope Estimator



Figure E255 Cold River: Aluminum Total



Figure E256 Cold River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.43
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 21.43
Adjusted Kruskal-Wallis statistic (H') = 21.43



Figure E257 Cold River: Arsenic Dissolved



Figure E258 Cold River: Arsenic Dissolved



Figure E259 Cold River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.188
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E260 Cold River: Arsenic Total

Seasonal Kendall



Figure E261 Cold River: Arsenic Total



Figure E262 Cold River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.845
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E263 Cold River: Barium Dissolved



Figure E264 Cold River: Barium Dissolved



Figure E265 Cold River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.82
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 15.82
Adjusted Kruskal-Wallis statistic (H') = 15.82



Figure E266 Cold River: Barium Total

Seasonal Kendall



Figure E267 Cold River: Barium Total



Figure E268 Cold River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.05804
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.05804
Adjusted Kruskal-Wallis statistic (H') = 0.05804



Figure E269 Cold River: Boron Dissolved



Figure E270 Cold River: Boron Dissolved



Figure E271 Cold River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.971
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E272 Cold River: Boron Total

Seasonal Kendall



Figure E273 Cold River: Boron Total



Figure E274 Cold River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.041
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.041
Adjusted Kruskal-Wallis statistic (H') = 2.041



Figure E275 Cold River: Cadmium Dissolved



Figure E276 Cold River: Cadmium Dissolved



Figure E277 Cold River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.39
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E278 Cold River: Cadmium Total

Seasonal Kendall



Figure E279 Cold River: Cadmium Total



Figure E280 Cold River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.505
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.505
Adjusted Kruskal-Wallis statistic (H') = 2.505



Figure E281 Cold River: Chromium Dissolved



Figure E282 Cold River: Chromium Dissolved



Figure E283 Cold River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.272
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 5 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 6.271
Adjusted Kruskal-Wallis statistic (H') = 6.272



Figure E284 Cold River: Chromium Total

Seasonal Kendall



Figure E285 Cold River: Chromium Total

Time Series



Figure E286 Cold River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.58
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.58
Adjusted Kruskal-Wallis statistic (H') = 4.58



Figure E287 Cold River: Cobalt Dissolved



Figure E288 Cold River: Cobalt Dissolved



Figure E289 Cold River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2628
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.2628
Adjusted Kruskal-Wallis statistic (H') = 0.2628



Figure E290 Cold River: Cobalt Total

Sen's Slope Estimator



Figure E291 Cold River: Cobalt Total

Time Series



Figure E292 Cold River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8754
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8754
Adjusted Kruskal-Wallis statistic (H') = 0.8754



Figure E293 Cold River: Copper Dissolved



Figure E294 Cold River: Copper Dissolved



Figure E295 Cold River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.104
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 6.104
Adjusted Kruskal-Wallis statistic (H') = 6.104



Figure E296 Cold River: Copper Total

Seasonal Kendall



Figure E297 Cold River: Copper Total



Figure E298 Cold River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8035
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E299 Cold River: Iron Dissolved



Figure E300 Cold River: Iron Dissolved



Figure E301 Cold River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.9071
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.9071
Adjusted Kruskal-Wallis statistic (H') = 0.9071



Figure E302 Cold River: Iron Total

Sen's Slope Estimator



Figure E303 Cold River: Iron Total



Figure E304 Cold River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01962
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E305 Cold River: Lithium Dissolved



Figure E306 Cold River: Lithium Dissolved



Figure E307 Cold River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8859
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E308 Cold River: Lithium Total

Sen's Slope Estimator



Figure E309 Cold River: Lithium Total



Figure E310 Cold River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.9723
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E311 Cold River: Manganese Dissolved



Figure E312 Cold River: Manganese Dissolved



Figure E313 Cold River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.0166
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E314 Cold River: Manganese Total

Sen's Slope Estimator



Figure E315 Cold River: Manganese Total



Figure E316 Cold River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.3014
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E317 Cold River: Molybdenum Dissolved



Figure E318 Cold River: Molybdenum Dissolved



Figure E319 Cold River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.7157
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E320 Cold River: Molybdenum Total

Sen's Slope Estimator



Figure E321 Cold River: Molybdenum Total



Figure E322 Cold River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.017
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.017
Adjusted Kruskal-Wallis statistic (H') = 1.017



Figure E323 Cold River: Nickel Dissolved



Figure E324 Cold River: Nickel Dissolved



Figure E325 Cold River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.262
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 4 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.262
Adjusted Kruskal-Wallis statistic (H') = 1.262



Figure E326 Cold River: Nickel Total

Sen's Slope Estimator



Figure E327 Cold River: Nickel Total

Time Series



Figure E328 Cold River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2059
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 6 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.2059
Adjusted Kruskal-Wallis statistic (H') = 0.2059



Figure E329 Cold River: Selenium Dissolved



Figure E330 Cold River: Selenium Dissolved



Figure E331 Cold River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.0005022
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 7 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.0005022
Adjusted Kruskal-Wallis statistic (H') = 0.0005022



Figure E332 Cold River: Selenium Total

Sen's Slope Estimator



Figure E333 Cold River: Selenium Total



Figure E334 Cold River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5102
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.5102
Adjusted Kruskal-Wallis statistic (H') = 0.5102



Figure E335 Cold River: Uranium Dissolved



Figure E336 Cold River: Uranium Dissolved



Figure E337 Cold River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.79
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.79
Adjusted Kruskal-Wallis statistic (H') = 3.79



Figure E338 Cold River: Uranium Total

Sen's Slope Estimator



Figure E339 Cold River: Uranium Total

Time Series



Figure E340 Cold River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01519
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.01519
Adjusted Kruskal-Wallis statistic (H') = 0.01519



Figure E341 Cold River: Vanadium Dissolved



Figure 342 Cold River: Vanadium Dissolved



Figure E343 Cold River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.867
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E344 Cold River: Vanadium Total

Seasonal Kendall



Figure E345 Cold River: Vanadium Total

Time Series



Figure E346 Cold River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.262
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.262
Adjusted Kruskal-Wallis statistic (H') = 4.262



Figure E347 Cold River: Zinc Dissolved



Figure E348 Cold River: Zinc Dissolved



Figure E349 North Saskatchewan River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 37.86
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E350 North Saskatchewan River: Aluminum Dissolved



Figure E351 North Saskatchewan River: Aluminum Dissolved



Figure E352 North Saskatchewan River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.936
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.936
Adjusted Kruskal-Wallis statistic (H') = 1.936



Figure E353 North Saskatchewan River: Aluminum Total

Sen's Slope Estimator



Figure E354 North Saskatchewan River: Aluminum Total



Figure E355 North Saskatchewan River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.08
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 5 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 29.08
Adjusted Kruskal-Wallis statistic (H') = 29.08



Figure E356 North Saskatchewan River: Arsenic Dissolved



Figure E357 North Saskatchewan River: Arsenic Dissolved



Figure E358 North Saskatchewan River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.991
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.991
Adjusted Kruskal-Wallis statistic (H') = 1.991



Figure E359 North Saskatchewan River: Arsenic Total

Sen's Slope Estimator



Figure E360 North Saskatchewan River: Arsenic Total



Figure E361 North Saskatchewan River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.7702
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 13 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.7702
Adjusted Kruskal-Wallis statistic (H') = 0.7702



Figure E362 North Saskatchewan River: Barium Dissolved



Figure E363 North Saskatchewan River: Barium Dissolved



Figure E364 North Saskatchewan River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 23.39
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 23.39
Adjusted Kruskal-Wallis statistic (H') = 23.39



Figure E365 North Saskatchewan River: Barium Total

Seasonal Kendall



Figure E366 North Saskatchewan River: Barium Total



Figure E367 North Saskatchewan River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8635
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 13 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8635
Adjusted Kruskal-Wallis statistic (H') = 0.8635



Figure E368 North Saskatchewan River: Beryllium Dissolved



Figure E369 North Saskatchewan River: Beryllium Dissolved



Figure E370 North Saskatchewan River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.08505
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 5 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.08505
Adjusted Kruskal-Wallis statistic (H') = 0.08505



Figure E371 North Saskatchewan River: Beryllium Total

Sen's Slope Estimator



Figure E372 North Saskatchewan River: Beryllium Total



Figure E373 North Saskatchewan River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.455
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.455
Adjusted Kruskal-Wallis statistic (H') = 1.455



Figure E374 North Saskatchewan River: Boron Dissolved



Figure E375 North Saskatchewan River: Boron Dissolved



Figure E376 North Saskatchewan River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.198
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.198
Adjusted Kruskal-Wallis statistic (H') = 1.198



Figure E377 North Saskatchewan River: Boron Total

Sen's Slope



Figure E378 North Saskatchewan River: Boron Total

Time Series



Figure E379 North Saskatchewan River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.479
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E380 North Saskatchewan River: Cadmium Dissolved



Figure E381 North Saskatchewan River: Cadmium Dissolved



Figure E382 North Saskatchewan River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 13.31
Adjusted Kruskal-Wallis statistic (H') = 13.31



Figure E383 North Saskatchewan River: Cadmium Total

Seasonal Kendall



Figure E384 North Saskatchewan River: Cadmium Total



Figure E385 North Saskatchewan River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.23
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 6 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 17.23
Adjusted Kruskal-Wallis statistic (H') = 17.23



Figure E386 North Saskatchewan River: Chromium Dissolved



Figure E387 North Saskatchewan River: Chromium Dissolved



Figure E388 North Saskatchewan River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.332
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 7.332
Adjusted Kruskal-Wallis statistic (H') = 7.332



Figure E389 North Saskatchewan River: Chromium Total

Seasonal Kendall



Figure E390 North Saskatchewan River: Chromium Total



Figure E391 North Saskatchewan River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.27
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 21.27
Adjusted Kruskal-Wallis statistic (H') = 21.27



Figure E392 North Saskatchewan River: Cobalt Dissolved



Figure E393 North Saskatchewan River: Cobalt Dissolved



Figure E394 North Saskatchewan River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6531
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure 395 North Saskatchewan River: Cobalt Total

Sen's Slope Estimator



Figure E396 North Saskatchewan River: Cobalt Total



Figure E397 North Saskatchewan River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.528
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.528
Adjusted Kruskal-Wallis statistic (H') = 1.528



Figure E398 North Saskatchewan River: Copper Dissolved



Figure E399 North Saskatchewan River: Copper Dissolved



Figure E400 North Saskatchewan River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.289
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.289
Adjusted Kruskal-Wallis statistic (H') = 1.289



Figure E401 North Saskatchewan River: Copper Total

Sen's Slope Estimator



Figure E402 North Saskatchewan River: Copper Total



Figure E403 North Saskatchewan River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.62
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E404 North Saskatchewan River: Iron Dissolved



Figure E405 North Saskatchewan River: Iron Dissolved



Figure E406 North Saskatchewan River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.687
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E407 North Saskatchewan River: Iron Total

Sen's Slope Estimator



Figure E408 North Saskatchewan River: Iron Total



Figure E409 North Saskatchewan River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.41
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 13.41
Adjusted Kruskal-Wallis statistic (H') = 13.41



Figure E410 North Saskatchewan River: Lead Dissolved



Figure E411 North Saskatchewan River: Lead Dissolved



Figure E412 North Saskatchewan River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.39
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E413 North Saskatchewan River: Lead Total

Sen's Slope Estimator



Figure E414 North Saskatchewan River: Lead Total



Figure E415 North Saskatchewan River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.661
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E416 North Saskatchewan River: Lithium Dissolved



Figure E417 North Saskatchewan River: Lithium Dissolved



Figure E418 North Saskatchewan River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.694
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E419 North Saskatchewan River: Lithium Total

Sen's Slope Estimator



Figure E420 North Saskatchewan River: Lithium Total



Figure E421 North Saskatchewan River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.001354
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E422 North Saskatchewan River: Manganese Dissolved



Figure E423 North Saskatchewan River: Manganese Dissolved



Figure E424 North Saskatchewan River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.573
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E425 North Saskatchewan River: Manganese Total

Sen's Slope Estimator



Figure E426 North Saskatchewan River: Manganese Total

Time Series



Figure E427 North Saskatchewan River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.04874
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.04874
Adjusted Kruskal-Wallis statistic (H') = 0.04874



Figure E428 North Saskatchewan River: Molybdenum Dissolved



Figure E429 North Saskatchewan River: Molybdenum Dissolved



Figure E430 North Saskatchewan River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1938
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.1938
Adjusted Kruskal-Wallis statistic (H') = 0.1938



Figure E431 North Saskatchewan River: Molybdenum Total

Sen's Slope Estimator



Figure E432 North Saskatchewan River: Molybdenum Total



Figure E433 North Saskatchewan River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5638
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.5638
Adjusted Kruskal-Wallis statistic (H') = 0.5638



Figure E434 North Saskatchewan River: Nickel Dissolved



Figure E435 North Saskatchewan River: Nickel Dissolved



Figure E436 North Saskatchewan River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.543
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.543
Adjusted Kruskal-Wallis statistic (H') = 4.543



Figure E437 North Saskatchewan River: Nickel Total

Seasonal Kendall



Figure E438 North Saskatchewan River: Nickel Total



Figure E439 North Saskatchewan River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8237
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8237
Adjusted Kruskal-Wallis statistic (H') = 0.8237



Figure E440 North Saskatchewan River: Selenium Dissolved



Figure E441 North Saskatchewan River: Selenium Dissolved



Figure E442 North Saskatchewan River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.359
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 3 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.359
Adjusted Kruskal-Wallis statistic (H') = 2.359



Figure E443 North Saskatchewan River: Selenium Total

Sen's Slope Estimator



Figure E444 North Saskatchewan River: Selenium Total



Figure E445 North Saskatchewan River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.09824
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 13 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.09824
Adjusted Kruskal-Wallis statistic (H') = 0.09824



Figure E446 North Saskatchewan River: Silver Total



Figure E447 North Saskatchewan River: Silver Total



Figure E448 North Saskatchewan River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.412
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 10 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.412
Adjusted Kruskal-Wallis statistic (H') = 1.412



Figure E449 North Saskatchewan River: Thallium Dissolved

Sen's Slope Estimator



Figure E450 North Saskatchewan River: Thallium Dissolved



Figure E451 North Saskatchewan River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.3402
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 8 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.3402
Adjusted Kruskal-Wallis statistic (H') = 0.3402



Figure E452 North Saskatchewan River: Thallium Total



Figure E453 North Saskatchewan River: Thallium Total



Figure E454 North Saskatchewan River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.512
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E455 North Saskatchewan River: Uranium Dissolved

Sen's Slope Estimator



Figure E456 North Saskatchewan River: Uranium Dissolved



Figure E457 North Saskatchewan River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 20.19
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E458 North Saskatchewan River: Uranium Total



Figure E459 North Saskatchewan River: Uranium Total



Figure E460 North Saskatchewan River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 32.25
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E461 North Saskatchewan River: Vanadium Dissolved

Seasonal Kendall



Figure E462 North Saskatchewan River: Vanadium Dissolved

Time Series



Figure E463 North Saskatchewan River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6013
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.6013
Adjusted Kruskal-Wallis statistic (H') = 0.6013



Figure E464 North Saskatchewan River: Vanadium Total



Figure E465 North Saskatchewan River: Vanadium Total



Figure E466 North Saskatchewan River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.36
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 21.36
Adjusted Kruskal-Wallis statistic (H') = 21.36



Figure E467 North Saskatchewan River: Zinc Dissolved

Seasonal Kendall



Figure E468 North Saskatchewan River: Zinc Dissolved

Time Series



Figure E469 North Saskatchewan River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.87
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 14.87
Adjusted Kruskal-Wallis statistic (H') = 14.87



Figure E470 North Saskatchewan River: Zinc Total



Figure E471 North Saskatchewan River: Zinc Total



Figure E472 Red Deer River (AB-SK): Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.02685
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E473 Red Deer River (AB-SK): Aluminum Dissolved



Figure E474 Red Deer River (AB-SK): Aluminum Dissolved



Figure E475 Red Deer River (AB-SK): Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.626
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E476 Red Deer River (AB-SK): Aluminum Total

Seasonal Kendall



Figure E477 Red Deer River (AB-SK): Aluminum Total



Figure E478 Red Deer River (AB-SK): Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.492
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E479 Red Deer River (AB-SK): Arsenic Dissolved



Figure E480 Red Deer River (AB-SK): Arsenic Dissolved



Figure E481 Red Deer River (AB-SK): Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.392
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.392
Adjusted Kruskal-Wallis statistic (H') = 1.392



Figure E482 Red Deer River (AB-SK): Arsenic Total

Sen's Slope Estimator



Figure E483 Red Deer River (AB-SK): Arsenic Total



Figure E484 Red Deer River (AB-SK): Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.91
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E485 Red Deer River (AB-SK): Barium Dissolved



Figure E486 Red Deer River (AB-SK): Barium Dissolved



Figure E487 Red Deer River (AB-SK): Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.17
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E488 Red Deer River (AB-SK): Barium Total

Seasonal Kendall



Figure E489 Red Deer River (AB-SK): Barium Total



Figure E490 Red Deer River (AB-SK): Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 10 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 5.12
Adjusted Kruskal-Wallis statistic (H') = 5.12



Figure E491 Red Deer River (AB-SK): Beryllium Dissolved



Figure E492 Red Deer River (AB-SK): Beryllium Dissolved



Figure E493 Red Deer River (AB-SK): Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.968
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 9.968
Adjusted Kruskal-Wallis statistic (H') = 9.968



Figure E494 Red Deer River (AB-SK): Beryllium Total

Seasonal Kendall



Figure E495 Red Deer River (AB-SK): Beryllium Total



Figure E496 Red Deer River (AB-SK): Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1261
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E497 Red Deer River (AB-SK): Boron Dissolved



Figure E498 Red Deer River (AB-SK): Boron Dissolved



Figure E499 Red Deer River (AB-SK): Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.000007459
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E500 Red Deer River (AB-SK): Boron Total

Sen's Slope Estimator



Figure E501 Red Deer River (AB-SK): Boron Total



Figure E502 Red Deer River (AB-SK): Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.739
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E503 Red Deer River (AB-SK): Cadmium Dissolved



Figure E504 Red Deer River (AB-SK): Cadmium Dissolved



Figure E505 Red Deer River (AB-SK): Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.03348
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E506 Red Deer River (AB-SK): Cadmium Total

Sen's Slope Estimator



Figure E507 Red Deer River (AB-SK): Cadmium Total



Figure E508 Red Deer River (AB-SK): Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.279
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.279
Adjusted Kruskal-Wallis statistic (H') = 3.279



Figure E509 Red Deer River (AB-SK): Chromium Dissolved



Figure E510 Red Deer River (AB-SK): Chromium Dissolved



Figure E511 Red Deer River (AB-SK): Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.479
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E512 Red Deer River (AB-SK): Chromium Total

Sen's Slope Estimator



Figure E513 Red Deer River (AB-SK): Chromium Total



Figure E514 Red Deer River (AB-SK): Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.167
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.167
Adjusted Kruskal-Wallis statistic (H') = 1.167



Figure 515 Red Deer River (AB-SK): Cobalt Dissolved



Figure E516 Red Deer River (AB-SK): Cobalt Dissolved



Figure 517 Red Deer River (AB-SK): Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.052
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E518 Red Deer River (AB-SK): Cobalt Total

Seasonal Kendall



Figure E519 Red Deer River (AB-SK): Cobalt Total



Figure E520 Red Deer River (AB-SK): Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.01
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E521 Red Deer River (AB-SK): Copper Dissolved



Figure E522 Red Deer River (AB-SK): Copper Dissolved



Figure E523 Red Deer River (AB-SK): Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8471
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E524 Red Deer River (AB-SK): Copper Total

Sen's Slope Estimator



Figure E525 Red Deer River (AB-SK): Copper Total



Figure E526 Red Deer River (AB-SK): Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.89
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 15.89
Adjusted Kruskal-Wallis statistic (H') = 15.89



Figure E527 Red Deer River (AB-SK): Iron Dissolved



Figure E528 Red Deer River (AB-SK): Iron Dissolved



Figure E529 Red Deer River (AB-SK): Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.03
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E530 Red Deer River (AB-SK): Iron Total

Seasonal Kendall



Figure E531 Red Deer River (AB-SK): Iron Total



Figure E532 Red Deer River (AB-SK): Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 19.41
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E533 Red Deer River (AB-SK): Lead Dissolved



Figure E534 Red Deer River (AB-SK): Lead Dissolved



Figure E535 Red Deer River (AB-SK): Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.975
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E536 Red Deer River (AB-SK): Lead Total

Seasonal Kendall



Figure E537 Red Deer River (AB-SK): Lead Total

Time Series



Figure E538 Red Deer River (AB-SK): Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.02017
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E539 Red Deer River (AB-SK): Lithium Dissolved



Figure E540 Red Deer River (AB-SK): Lithium Dissolved



Figure E541 Red Deer River (AB-SK): Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5478
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E542 Red Deer River (AB-SK): Lithium Total

Sen's Slope Estimator



Figure E543 Red Deer River (AB-SK): Lithium Total



Figure E544 Red Deer River (AB-SK): Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 12.61
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E545 Red Deer River (AB-SK): Manganese Dissolved



Figure E546 Red Deer River (AB-SK): Manganese Dissolved



Figure E547 Red Deer River (AB-SK): Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 25.47
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E548 Red Deer River (AB-SK): Manganese Total

Seasonal Kendall



Figure E549 Red Deer River (AB-SK): Manganese Total



Figure E550 Red Deer River (AB-SK): Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.848
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 9.848
Adjusted Kruskal-Wallis statistic (H') = 9.848



Figure E551 Red Deer River (AB-SK): Molybdenum Dissolved



Figure E552 Red Deer River (AB-SK): Molybdenum Dissolved



Figure E553 Red Deer River (AB-SK): Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.453
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E554 Red Deer River (AB-SK): Molybdenum Total

Seasonal Kendall



Figure E555 Red Deer River (AB-SK): Molybdenum Total



Figure E556 Red Deer River (AB-SK): Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5081
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E557 Red Deer River (AB-SK): Nickel Dissolved



Figure E558 Red Deer River (AB-SK): Nickel Dissolved



Figure E559 Red Deer River (AB-SK): Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.7122
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E560 Red Deer River (AB-SK): Nickel Total

Sen's Slope Estimator



Figure E561 Red Deer River (AB-SK): Nickel Total

Time Series



Figure E562 Red Deer River (AB-SK): Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.985
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 8.985
Adjusted Kruskal-Wallis statistic (H') = 8.985



Figure E563 Red Deer River (AB-SK): Selenium Dissolved



Figure E564 Red Deer River (AB-SK): Selenium Dissolved



Figure E565 Red Deer River (AB-SK): Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.74
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E566 Red Deer River (AB-SK): Selenium Total

Seasonal Kendall



Figure E567 Red Deer River (AB-SK): Selenium Total



Figure E568 Red Deer River (AB-SK): Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.581
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 5.581
Adjusted Kruskal-Wallis statistic (H') = 5.581



Figure E569 Red Deer River (AB-SK): Silver Total



Figure E570 Red Deer River (AB-SK): Silver Total



Figure E571 Red Deer River (AB-SK): Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.056
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 2.056
Adjusted Kruskal-Wallis statistic (H') = 2.056



Figure E572 Red Deer River (AB-SK): Thallium Dissolved

Sen's Slope Estimator



Figure E573 Red Deer River (AB-SK): Thallium Dissolved



Figure E574 Red Deer River (AB-SK): Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.132
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E575 Red Deer River (AB-SK): Thallium Total



Figure E576 Red Deer River (AB-SK): Thallium Total



Figure E577 Red Deer River (AB-SK): Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.524
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.524
Adjusted Kruskal-Wallis statistic (H') = 1.524



Figure E578 Red Deer River (AB-SK): Uranium Dissolved

Sen's Slope Estimator



Figure E579 Red Deer River (AB-SK): Uranium Dissolved



Figure E580 Red Deer River (AB-SK): Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.979
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E581 Red Deer River (AB-SK): Uranium Total



Figure E582 Red Deer River (AB-SK): Uranium Total



Figure E583 Red Deer River (AB-SK): Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 33.27
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E584 Red Deer River (AB-SK): Vanadium Dissolved

Seasonal Kendall



Figure E585 Red Deer River (AB-SK): Vanadium Dissolved



Figure E586 Red Deer River (AB-SK): Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.224
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E587 Red Deer River (AB-SK): Vanadium Total



Figure E588 Red Deer River (AB-SK): Vanadium Total



Figure E589 Red Deer River (AB-SK): Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 22.04
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E590 Red Deer River (AB-SK): Zinc Dissolved

Seasonal Kendall



Figure E591 Red Deer River (AB-SK): Zinc Dissolved



Figure E592 Red Deer River (AB-SK): Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4296
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E593 Red Deer River (AB-SK): Zinc Total



Figure E594 Red Deer River (AB-SK): Zinc Total

Time Series



Figure E595 South Saskatchewan River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 25.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E596 South Saskatchewan River: Aluminum Dissolved



Figure E597 South Saskatchewan River: Aluminum Dissolved



Figure E598 South Saskatchewan River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.558
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E599 South Saskatchewan River: Aluminum Total

Seasonal Kendall



Figure E600 South Saskatchewan River: Aluminum Total



Figure E601 South Saskatchewan River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 76.82
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E602 South Saskatchewan River: Arsenic Dissolved



Figure E603 South Saskatchewan River: Arsenic Dissolved



Figure E604 South Saskatchewan River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 47.07
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E605 South Saskatchewan River: Arsenic Total

Seasonal Kendall



Figure E606 South Saskatchewan River: Arsenic Total



Figure E607 South Saskatchewan River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1243
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E608 South Saskatchewan River: Barium Dissolved



Figure E609 South Saskatchewan River: Barium Dissolved



Figure E610 South Saskatchewan River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01694
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E611 South Saskatchewan River: Barium Total

Sen's Slope Estimator



Figure E612 South Saskatchewan River: Barium Total



Figure E613 South Saskatchewan River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.15
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E614 South Saskatchewan River: Beryllium Total



Figure E615 South Saskatchewan River: Beryllium Total



Figure E616 South Saskatchewan River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.555
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E617 South Saskatchewan River: Boron Dissolved

Seasonal Kendall



Figure E618 South Saskatchewan River: Boron Dissolved



Figure E619 South Saskatchewan River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.636
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E620 South Saskatchewan River: Boron Total



Figure E621 South Saskatchewan River: Boron Total



Figure E622 South Saskatchewan River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.276
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 5.276
Adjusted Kruskal-Wallis statistic (H') = 5.276



Figure E623 South Saskatchewan River: Cadmium Dissolved

Seasonal Kendall



Figure E624 South Saskatchewan River: Cadmium Dissolved



Figure E625 South Saskatchewan River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.02919
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E626 South Saskatchewan River: Cadmium Total



Figure E627 South Saskatchewan River: Cadmium Total



Figure E628 South Saskatchewan River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.38
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E629 South Saskatchewan River: Chromium Dissolved

Seasonal Kendall



Figure E630 South Saskatchewan River: Chromium Dissolved

Time Series



Figure E631 South Saskatchewan River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.03552
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.03552
Adjusted Kruskal-Wallis statistic (H') = 0.03552



Figure E632 South Saskatchewan River: Chromium Total



Figure E633 South Saskatchewan River: Chromium Total



Figure E634 South Saskatchewan River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.33
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E635 South Saskatchewan River: Cobalt Dissolved

Seasonal Kendall



Figure E636 South Saskatchewan River: Cobalt Dissolved

Time Series



Figure E637 South Saskatchewan River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 22.99
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 22.99
Adjusted Kruskal-Wallis statistic (H') = 22.99



Figure E638 South Saskatchewan River: Cobalt Total



Figure E639 South Saskatchewan River: Cobalt Total



Figure E640 South Saskatchewan River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.49
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E641 South Saskatchewan River: Copper Dissolved

Seasonal Kendall



Figure E642 South Saskatchewan River: Copper Dissolved



Figure E643 South Saskatchewan River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.71
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E644 South Saskatchewan River: Copper Total



Figure E645 South Saskatchewan River: Copper Total



Figure E646 South Saskatchewan River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.854
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E647 South Saskatchewan River: Iron Dissolved

Sen's Slope Estimator



Figure E648 South Saskatchewan River: Iron Dissolved



Figure E649 South Saskatchewan River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.209
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E650 South Saskatchewan River: Iron Total



Figure E651 South Saskatchewan River: Iron Total



Figure E652 South Saskatchewan River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 23.14
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E653 South Saskatchewan River: Lead Dissolved

Seasonal Kendall



Figure E654 South Saskatchewan River: Lead Dissolved

Time Series



Figure E655 South Saskatchewan River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.682
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E656 South Saskatchewan River: Lead Total



Figure E657 South Saskatchewan River: Lead Total



Figure E658 South Saskatchewan River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 22.6
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E659 South Saskatchewan River: Lithium Dissolved

Seasonal Kendall



Figure E660 South Saskatchewan River: Lithium Dissolved



Figure E661 South Saskatchewan River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.78
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E662 South Saskatchewan River: Lithium Total



Figure E663 South Saskatchewan River: Lithium Total



Figure E664 South Saskatchewan River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.326
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E665 South Saskatchewan River: Manganese Dissolved

Seasonal Kendall



Figure E666 South Saskatchewan River: Manganese Dissolved



Figure E667 South Saskatchewan River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 43.28
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E668 South Saskatchewan River: Manganese Total



Figure E669 South Saskatchewan River: Manganese Total



Figure E670 South Saskatchewan River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.0007354
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E671 South Saskatchewan River: Molybdenum Dissolved

Sen's Slope Estimator



Figure E672 South Saskatchewan River: Molybdenum Dissolved



Figure E673 South Saskatchewan River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.773
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E674 South Saskatchewan River: Molybdenum Total



Figure E675 South Saskatchewan River: Molybdenum Total



Figure E676 South Saskatchewan River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.92
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E677 South Saskatchewan River: Nickel Dissolved

Seasonal Kendall



Figure E678 South Saskatchewan River: Nickel Dissolved



Figure E679 South Saskatchewan River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.73
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E680 South Saskatchewan River: Nickel Total



Figure E681 South Saskatchewan River: Nickel Total



Figure E682 South Saskatchewan River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 48.54
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E683 South Saskatchewan River: Selenium Dissolved

Seasonal Kendall



Figure E684 South Saskatchewan River: Selenium Dissolved

Time Series



Figure E685 South Saskatchewan River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 50.52
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E686 South Saskatchewan River: Selenium Total



Figure E687 South Saskatchewan River: Selenium Total



Figure E688 South Saskatchewan River: Silver Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8155
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.8155
Adjusted Kruskal-Wallis statistic (H') = 0.8155



Figure E689 South Saskatchewan River: Silver Dissolved

Sen's Slope Estimator



Figure E690 South Saskatchewan River: Silver Dissolved



Figure E691 South Saskatchewan River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.209
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E692 South Saskatchewan River: Silver Total



Figure E693 South Saskatchewan River: Silver Total



Figure E694 South Saskatchewan River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.341
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E695 South Saskatchewan River: Thallium Dissolved

Sen's Slope Estimator



Figure E696 South Saskatchewan River: Thallium Dissolved



Figure E697 South Saskatchewan River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.863
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E698 South Saskatchewan River: Thallium Total



Figure E699 South Saskatchewan River: Thallium Total



Figure E700 South Saskatchewan River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.369
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E701 South Saskatchewan River: Uranium Dissolved

Sen's Slope Estimator



Figure E702 South Saskatchewan River: Uranium Dissolved



Figure E703 South Saskatchewan River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1077
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E704 South Saskatchewan River: Uranium Total



Figure E705 South Saskatchewan River: Uranium Total



Figure E706 South Saskatchewan River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 79.1
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 79.1
Adjusted Kruskal-Wallis statistic (H') = 79.1



Figure E707 South Saskatchewan River: Vanadium Dissolved

Seasonal Kendall



Figure E708 South Saskatchewan River: Vanadium Dissolved



Figure E709 South Saskatchewan River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 20.22
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E710 South Saskatchewan River: Vanadium Total



Figure E711 South Saskatchewan River: Vanadium Total



Figure E712 South Saskatchewan River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 47.19
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E713 South Saskatchewan River: Zinc Dissolved

Seasonal Kendall



Figure E714 South Saskatchewan River: Zinc Dissolved



Figure E715 South Saskatchewan River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.658
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E716 South Saskatchewan River: Zinc Total



Figure E717 South Saskatchewan River: Zinc Total

Appendix E: Metals Trending Graphs Part 2
Saskatchewan/Manitoba Border



Figure E709 Assiniboine River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5498
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.5498
Adjusted Kruskal-Wallis statistic (H') = 0.5498



Figure E710 Assiniboine River: Aluminum Dissolved



Figure E711 Assiniboine River: Aluminum Dissolved



Figure E712 Assiniboine River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 46.63
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E713 Assiniboine River: Aluminum Total

Seasonal Kendall



Figure E714 Assiniboine River: Aluminum Total



Figure E715 Assiniboine River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 103.1
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E716 Assiniboine River: Arsenic Dissolved



Figure E717 Assiniboine River: Arsenic Dissolved



Figure E718 Assiniboine River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 109.2
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E719 Assiniboine River: Arsenic Total

Seasonal Kendall



Figure E720 Assiniboine River: Arsenic Total

Time Series



Figure E721 Assiniboine River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.652
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E722 Assiniboine River: Barium Dissolved



Figure E723 Assiniboine River: Barium Dissolved



Figure E724 Assiniboine River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1205
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E725 Assiniboine River: Barium Total

Sen's Slope Estimator



Figure E726 Assiniboine River: Barium Total



Figure E727 Assiniboine River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.744
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 5.744
Adjusted Kruskal-Wallis statistic (H') = 5.744



Figure E728 Assiniboine River: Beryllium Dissolved



Figure E729 Assiniboine River: Beryllium Dissolved



Figure E730 Assiniboine River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.63
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E731 Assiniboine River: Beryllium Total

Seasonal Kendall



Figure E732 Assiniboine River: Beryllium Total

Time Series



Figure E733 Assiniboine River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.114
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E734 Assiniboine River: Boron Dissolved



Figure E735 Assiniboine River: Boron Dissolved



Figure E736 Assiniboine River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.76
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E737 Assiniboine River: Boron Total

Seasonal Kendall



Figure E738 Assiniboine River: Boron Total



Figure E739 Assiniboine River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.846
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.846
Adjusted Kruskal-Wallis statistic (H') = 1.846



Figure E740 Assiniboine River: Cadmium Dissolved



Figure E741 Assiniboine River: Cadmium Dissolved



Figure E742 Assiniboine River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 12.63
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 12.63
Adjusted Kruskal-Wallis statistic (H') = 12.63



Figure E743 Assiniboine River: Cadmium Total

Seasonal Kendall



Figure E744 Assiniboine River: Cadmium Total



Figure E745 Assiniboine River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.728
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.728
Adjusted Kruskal-Wallis statistic (H') = 3.728



Figure E746 Assiniboine River: Chromium Dissolved



Figure E747 Assiniboine River: Chromium Dissolved



Figure E748 Assiniboine River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 30.58
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E749 Assiniboine River: Chromium Total

Seasonal Kendall



Figure E750 Assiniboine River: Chromium Total



Figure E751 Assiniboine River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 24.17
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E752 Assiniboine River: Cobalt Dissolved



Figure E753 Assiniboine River: Cobalt Dissolved



Figure E754 Assiniboine River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 56.59
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E755 Assiniboine River: Cobalt Total

Seasonal Kendall



Figure E756 Assiniboine River: Cobalt Total



Figure E757 Assiniboine River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 24.88
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 24.88
Adjusted Kruskal-Wallis statistic (H') = 24.88



Figure E758 Assiniboine River: Copper Dissolved



Figure E759 Assiniboine River: Copper Dissolved



Figure E760 Assiniboine River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.34
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E761 Assiniboine River: Copper Total

Seasonal Kendall



Figure E762 Assiniboine River: Copper Total

Time Series



Figure E763 Assiniboine River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 69.53
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E764 Assiniboine River: Iron Dissolved



Figure E765 Assiniboine River: Iron Dissolved



Figure E766 Assiniboine River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.05
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E767 Assiniboine River: Iron Total

Seasonal Kendall



Figure E768 Assiniboine River: Iron Total



Figure E769 Assiniboine River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.167
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E770 Assiniboine River: Lead Dissolved



Figure E771 Assiniboine River: Lead Dissolved



Figure E772 Assiniboine River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 53.37
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E773 Assiniboine River: Lead Total

Seasonal Kendall



Figure E774 Assiniboine River: Lead Total

Time Series



Figure E775 Assiniboine River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.314
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E776 Assiniboine River: Lithium Dissolved



Figure E777 Assiniboine River: Lithium Dissolved



Figure E778 Assiniboine River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.787
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E779 Assiniboine River: Lithium Total

Sen's Slope Estimator



Figure E780 Assiniboine River: Lithium Total

Time Series



Figure E781 Assiniboine River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.3
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E782 Assiniboine River: Manganese Dissolved



Figure E783 Assiniboine River: Manganese Dissolved



Figure E784 Assiniboine River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.02474
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E785 Assiniboine River: Manganese Total

Sen's Slope Estimator



Figure E786 Assiniboine River: Manganese Total



Figure E787 Assiniboine River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.32
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E788 Assiniboine River: Molybdenum Dissolved



Figure E789 Assiniboine River: Molybdenum Dissolved



Figure E790 Assiniboine River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.82
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E791 Assiniboine River: Molybdenum Total

Seasonal Kendall



Figure E792 Assiniboine River: Molybdenum Total



Figure E793 Assiniboine River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.19
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E794 Assiniboine River: Nickel Dissolved



Figure E795 Assiniboine River: Nickel Dissolved



Figure E796 Assiniboine River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 58.44
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E797 Assiniboine River: Nickel Total

Seasonal Kendall



Figure E798 Assiniboine River: Nickel Total



Figure E799 Assiniboine River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.1
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E800 Assiniboine River: Selenium Dissolved



Figure E801 Assiniboine River: Selenium Dissolved



Figure E802 Assiniboine River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.27
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E803 Assiniboine River: Selenium Total

Seasonal Kendall



Figure E804 Assiniboine River: Selenium Total



Figure E805 Assiniboine River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.18
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 29.18
Adjusted Kruskal-Wallis statistic (H') = 29.18



Figure E806 Assiniboine River: Silver Total



Figure E807 Assiniboine River: Silver Total



Figure E808 Assiniboine River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.103
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.103
Adjusted Kruskal-Wallis statistic (H') = 1.103



Figure E809 Assiniboine River: Thallium Dissolved

Sen's Slope Estimator



Figure E810 Assiniboine River: Thallium Dissolved



Figure E811 Assiniboine River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 24.82
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E812 Assiniboine River: Thallium Total



Figure E813 Assiniboine River: Thallium Total



Figure E814 Assiniboine River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.318
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E815 Assiniboine River: Uranium Dissolved

Seasonal Kendall



Figure E816 Assiniboine River: Uranium Dissolved



Figure E817 Assiniboine River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.046
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E818 Assiniboine River: Uranium Total



Figure E819 Assiniboine River: Uranium Total



Figure E820 Assiniboine River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 112
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E821 Assiniboine River: Vanadium Dissolved

Seasonal Kendall



Figure E822 Assiniboine River: Vanadium Dissolved



Figure E823 Assiniboine River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 100.2
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E824 Assiniboine River: Vanadium Total



Figure E825 Assiniboine River: Vanadium Total



Figure E826 Assiniboine River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 38.3
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E827 Assiniboine River: Zinc Dissolved

Seasonal Kendall



Figure E828 Assiniboine River: Zinc Dissolved

Time Series



Figure E829 Assiniboine River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.562
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E830 Assiniboine River: Zinc Total



Figure E831 Assiniboine River: Zinc Total

Time Series



Figure E832 Carrot River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.08
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E833 Carrot River: Aluminum Dissolved



Figure E834 Carrot River: Aluminum Dissolved



Figure E835 Carrot River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 62.36
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E836 Carrot River: Aluminum Total

Seasonal Kendall



Figure E837 Carrot River: Aluminum Total

Time Series



Figure E838 Carrot River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.595
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 38 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 8.594
Adjusted Kruskal-Wallis statistic (H') = 8.595



Figure E839 Carrot River: Arsenic Dissolved



Figure E840 Carrot River: Arsenic Dissolved



Figure E841 Carrot River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.736
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E842 Carrot River: Arsenic Total

Seasonal Kendall



Figure E843 Carrot River: Arsenic Total

Time Series



Figure E844 Carrot River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 60.07
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E845 Carrot River: Barium Dissolved



Figure E846 Carrot River: Barium Dissolved



Figure E847 Carrot River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 40.74
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E848 Carrot River: Barium Total

Seasonal Kendall



Figure E849 Carrot River: Barium Total

Time Series



Figure E850 Carrot River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1211
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.1211
Adjusted Kruskal-Wallis statistic (H') = 0.1211



Figure E851 Carrot River: Beryllium Dissolved



Figure E852 Carrot River: Beryllium Dissolved



Figure E853 Carrot River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 59.19
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E854 Carrot River: Beryllium Total

Seasonal Kendall



Figure E855 Carrot River: Beryllium Total

Time Series



Figure E856 Carrot River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 35.43
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E857 Carrot River: Boron Dissolved



Figure E858 Carrot River: Boron Dissolved



Figure E859 Carrot River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.77
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E860 Carrot River: Boron Total

Seasonal Kendall



Figure E861 Carrot River: Boron Total

Time Series



Figure E862 Carrot River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.9148
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E863 Carrot River: Cadmium Dissolved



Figure E864 Carrot River: Cadmium Dissolved



Figure E865 Carrot River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.69
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 27.69
Adjusted Kruskal-Wallis statistic (H') = 27.69



Figure E866 Carrot River: Cadmium Total

Seasonal Kendall



Figure E867 Carrot River: Cadmium Total

Time Series



Figure E868 Carrot River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.622
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 4.622
Adjusted Kruskal-Wallis statistic (H') = 4.622



Figure E869 Carrot River: Chromium Dissolved



Figure E870 Carrot River: Chromium Dissolved



Figure E871 Carrot River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 42.55
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E872 Carrot River: Chromium Total

Seasonal Kendall



Figure E873 Carrot River: Chromium Total

Time Series



Figure E874 Carrot River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 43.66
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E875 Carrot River: Cobalt Dissolved



Figure E876 Carrot River: Cobalt Dissolved



Figure E877 Carrot River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.195
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.195
Adjusted Kruskal-Wallis statistic (H') = 3.195



Figure 878 Carrot River: Cobalt Total

Sen's Slope Estimator



Figure E879 Carrot River: Cobalt Total

Time Series



Figure E880 Carrot River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 23.45
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E881 Carrot River: Copper Dissolved



Figure E882 Carrot River: Copper Dissolved



Figure E883 Carrot River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.16
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E884 Carrot River: Copper Total

Seasonal Kendall



Figure E885 Carrot River: Copper Total



Figure E886 Carrot River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 24.51
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E887 Carrot River: Iron Dissolved



Figure E888 Carrot River: Iron Dissolved



Figure E889 Carrot River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.4
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E890 Carrot River: Iron Total

Seasonal Kendall



Figure E891 Carrot River: Iron Total

Time Series



Figure E892 Carrot River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.65
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E893 Carrot River: Lead Dissolved



Figure E894 Carrot River: Lead Dissolved



Figure E895 Carrot River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 56.7
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E896 Carrot River: Lead Total

Seasonal Kendall



Figure 897 Carrot River: Lead Total

Time Series



Figure E898 Carrot River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 59.66
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E899 Carrot River: Lithium Dissolved



Figure E900 Carrot River: Lithium Dissolved



Figure E901 Carrot River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 64.33
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E902 Carrot River: Lithium Total

Seasonal Kendall



Figure E903 Carrot River: Lithium Total



Figure E904 Carrot River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 73.75
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E905 Carrot River: Manganese Dissolved



Figure E906 Carrot River: Manganese Dissolved



Figure E907 Carrot River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 62.2
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E908 Carrot River: Manganese Total

Seasonal Kendall



Figure E909 Carrot River: Manganese Total



Figure E910 Carrot River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.444
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E911 Carrot River: Molybdenum Dissolved



Figure E912 Carrot River: Molybdenum Dissolved



Figure E913 Carrot River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8411
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E914 Carrot River: Molybdenum Total

Sen's Slope Estimator



Figure E915 Carrot River: Molybdenum Total



Figure E916 Carrot River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.483
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E917 Carrot River: Nickel Dissolved



Figure E918 Carrot River: Nickel Dissolved



Figure E919 Carrot River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.173
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E920 Carrot River: Nickel Total

Seasonal Kendall



Figure E921 Carrot River: Nickel Total

Time Series



Figure E922 Carrot River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6839
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.6839
Adjusted Kruskal-Wallis statistic (H') = 0.6839



Figure E923 Carrot River: Selenium Dissolved



Figure E924 Carrot River: Selenium Dissolved



Figure E925 Carrot River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6868
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.6868
Adjusted Kruskal-Wallis statistic (H') = 0.6868



Figure E926 Carrot River: Selenium Total

Sen's Slope Estimator



Figure E927 Carrot River: Selenium Total



Figure E928 Carrot River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 27.12
Adjusted Kruskal-Wallis statistic (H') = 27.12



Figure E929 Carrot River: Silver Total



Figure E930 Carrot River: Silver Total



Figure E931 Carrot River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.515
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 8.515
Adjusted Kruskal-Wallis statistic (H') = 8.515



Figure E932 Carrot River: Thallium Dissolved

Seasonal Kendall



Figure E933 Carrot River: Thallium Dissolved

Time Series



Figure E934 Carrot River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 38.52
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E935 Carrot River: Thallium Total



Figure E936 Carrot River: Thallium Total



Figure E937 Carrot River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 28.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E938 Carrot River: Uranium Dissolved

Seasonal Kendall



Figure E939 Carrot River: Uranium Dissolved

Time Series



Figure E940 Carrot River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 21.92
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E941 Carrot River: Uranium Total



Figure E942 Carrot River: Uranium Total



Figure E943 Carrot River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 62.39
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E944 Carrot River: Vanadium Dissolved

Seasonal Kendall



Figure E945 Carrot River: Vanadium Dissolved

Time Series



Figure E946 Carrot River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 76.93
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E947 Carrot River: Vanadium Total



Figure E948 Carrot River: Vanadium Total



Figure E949 Carrot River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.88
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E950 Carrot River: Zinc Dissolved

Seasonal Kendall



Figure E951 Carrot River: Zinc Dissolved

Time Series



Figure E952 Carrot River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.03
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 27.03
Adjusted Kruskal-Wallis statistic (H') = 27.03



Figure E953 Carrot River: Zinc Total



Figure E954 Carrot River: Zinc Total



Figure E955 Churchill River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 23.13
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E956 Churchill River: Aluminum Dissolved



Figure E957 Churchill River: Aluminum Dissolved



Figure E958 Churchill River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 26.95
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E959 Churchill River: Aluminum Total

Seasonal Kendall



Figure E960 Churchill River: Aluminum Total



Figure E961 Churchill River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.527
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E962 Churchill River: Arsenic Dissolved



Figure E963 Churchill River: Arsenic Dissolved



Figure E964 Churchill River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.441
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E965 Churchill River: Arsenic Total

Sen's Slope Estimator



Figure E966 Churchill River: Arsenic Total

Time Series



Figure E967 Churchill River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.201
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E968 Churchill River: Barium Dissolved



Figure E969 Churchill River: Barium Dissolved



Figure E970 Churchill River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.507
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E971 Churchill River: Barium Total

Sen's Slope Estimator



Figure E972 Churchill River: Barium Total



Figure E973 Churchill River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.578
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E974 Churchill River: Beryllium Dissolved



Figure E975 Churchill River: Beryllium Dissolved



Figure E976 Churchill River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.63
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E977 Churchill River: Beryllium Total

Seasonal Kendall



Figure E978 Churchill River: Beryllium Total

Time Series



Figure E979 Churchill River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.545
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E980 Churchill River: Boron Dissolved



Figure E981 Churchill River: Boron Dissolved



Figure E982 Churchill River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.09
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E983 Churchill River: Boron Total

Sen's Slope Estimator



Figure E984 Churchill River: Boron Total



Figure E985 Churchill River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.809
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E986 Churchill River: Cadmium Dissolved



Figure E987 Churchill River: Cadmium Dissolved



Figure E988 Churchill River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.002354
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E989 Churchill River: Cadmium Total

Sen's Slope Estimator



Figure E990 Churchill River: Cadmium Total

Time Series



Figure E991 Churchill River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.07018
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E992 Churchill River: Chromium Dissolved



Figure E993 Churchill River: Chromium Dissolved



Figure E994 Churchill River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.87
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E995 Churchill River: Chromium Total

Seasonal Kendall



Figure E996 Churchill River: Chromium Total

Time Series



Figure E997 Churchill River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.988
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E998 Churchill River: Cobalt Dissolved



Figure E999 Churchill River: Cobalt Dissolved



Figure E1000 Churchill River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 22.51
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1001 Churchill River: Cobalt Total

Seasonal Kendall



Figure E1002 Churchill River: Cobalt Total



Figure E1003 Churchill River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.018
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1004 Churchill River: Copper Dissolved



Figure E1005 Churchill River: Copper Dissolved



Figure E1006 Churchill River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.24
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1007 Churchill River: Copper Total

Seasonal Kendall



Figure E1008 Churchill River: Copper Total



Figure E1009 Churchill River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4801
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1010 Churchill River: Iron Dissolved



Figure E1011 Churchill River: Iron Dissolved



Figure E1012 Churchill River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 20.01
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1013 Churchill River: Iron Total

Seasonal Kendall



Figure E1014 Churchill River: Iron Total



Figure E1015 Churchill River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1346
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1016 Churchill River: Lead Dissolved



Figure E1017 Churchill River: Lead Dissolved



Figure E1018 Churchill River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.434
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1019 Churchill River: Lead Total

Seasonal Kendall



Figure E1020 Churchill River: Lead Total

Time Series



Figure E1021 Churchill River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.596
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1022 Churchill River: Lithium Dissolved



Figure E1023 Churchill River: Lithium Dissolved



Figure E1024 Churchill River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.95
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1025 Churchill River: Lithium Total

Sen's Slope Estimator



Figure E1026 Churchill River: Lithium Total

Time Series



Figure E1027 Churchill River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.68
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1028 Churchill River: Manganese Dissolved



Figure E1029 Churchill River: Manganese Dissolved



Figure E1030 Churchill River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1031 Churchill River: Manganese Total

Seasonal Kendall



Figure E1032 Churchill River: Manganese Total



Figure E1033 Churchill River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.003738
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1034 Churchill River: Molybdenum Dissolved



Figure E1035 Churchill River: Molybdenum Dissolved



Figure E1036 Churchill River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.04152
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1037 Churchill River: Molybdenum Total

Sen's Slope Estimator



Figure E1038 Churchill River: Molybdenum Total

Time Series



Figure E1039 Churchill River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.08
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1040 Churchill River: Nickel Dissolved



Figure E1041 Churchill River: Nickel Dissolved



Figure E1042 Churchill River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.98
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1043 Churchill River: Nickel Total

Seasonal Kendall



Figure E1044 Churchill River: Nickel Total

Time Series



Figure E1045 Churchill River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.446
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1046 Churchill River: Thallium Dissolved



Figure E1047 Churchill River: Thallium Dissolved



Figure E1048 Churchill River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 16.53
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1049 Churchill River: Thallium Total

Seasonal Kendall



Figure E1050 Churchill River: Thallium Total



Figure E1051 Churchill River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.61
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1052 Churchill River: Uranium Dissolved



Figure E1053 Churchill River: Uranium Dissolved



Figure E1054 Churchill River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 18.81
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1055 Churchill River: Uranium Total

Seasonal Kendall



Figure E1056 Churchill River: Uranium Total

Time Series



Figure E1057 Churchill River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 26.37
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1058 Churchill River: Vanadium Dissolved



Figure E1059 Churchill River: Vanadium Dissolved



Figure E1060 Churchill River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.42
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1061 Churchill River: Vanadium Total

Seasonal Kendall



Figure E1062 Churchill River: Vanadium Total

Time Series



Figure E1063 Churchill River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.018
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1064 Churchill River: Zinc Dissolved



Figure E1065 Churchill River: Zinc Dissolved



Figure E1066 Churchill River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01591
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1067 Churchill River: Zinc Total

Sen's Slope Estimator



Figure E1068 Churchill River: Zinc Total



Figure E1069 Qu'Appelle River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5997
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1070 Qu'Appelle River: Aluminum Dissolved



Figure E1071 Qu'Appelle River: Aluminum Dissolved



Figure E1072 Qu'Appelle River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1073 Qu'Appelle River: Aluminum Total

Seasonal Kendall



Figure E1074 Qu'Appelle River: Aluminum Total



Figure E1075 Qu'Appelle River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.404
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1076 Qu'Appelle River: Arsenic Dissolved



Figure E1077 Qu'Appelle River: Arsenic Dissolved



Figure E1078 Qu'Appelle River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 16.85
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1079 Qu'Appelle River: Arsenic Total

Seasonal Kendall



Figure E1080 Qu'Appelle River: Arsenic Total

Time Series



Figure E1081 Qu'Appelle River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.924
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1082 Qu'Appelle River: Barium Dissolved



Figure E1083 Qu'Appelle River: Barium Dissolved



Figure E1084 Qu'Appelle River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.4
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1085 Qu'Appelle River: Barium Total

Seasonal Kendall



Figure E1086 Qu'Appelle River: Barium Total

Time Series



Figure E1087 Qu'Appelle River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4082
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 0.4082
Adjusted Kruskal-Wallis statistic (H') = 0.4082



Figure E1088 Qu'Appelle River: Beryllium Dissolved



Figure E1089 Qu'Appelle River: Beryllium Dissolved



Figure E1090 Qu'Appelle River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 43.64
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1091 Qu'Appelle River: Beryllium Total

Seasonal Kendall



Figure E1092 Qu'Appelle River: Beryllium Total



Figure E1093 Qu'Appelle River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.668
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1094 Qu'Appelle River: Boron Dissolved



Figure E1095 Qu'Appelle River: Boron Dissolved



Figure E1096 Qu'Appelle River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.654
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1097 Qu'Appelle River: Boron Total

Sen's Slope Estimator



Figure E1098 Qu'Appelle River: Boron Total

Time Series



Figure E1099 Qu'Appelle River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1707
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1100 Qu'Appelle River: Cadmium Dissolved



Figure E1101 Qu'Appelle River: Cadmium Dissolved



Figure E1102 Qu'Appelle River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 28.66
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 28.66
Adjusted Kruskal-Wallis statistic (H') = 28.66



Figure E1103 Qu'Appelle River: Cadmium Total

Seasonal Kendall



Figure E1104 Qu'Appelle River: Cadmium Total

Time Series



Figure E1105 Qu'Appelle River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.8994
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1106 Qu'Appelle River: Chromium Dissolved



Figure E1107 Qu'Appelle River: Chromium Dissolved



Figure E1108 Qu'Appelle River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 32.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1109 Qu'Appelle River: Chromium Total

Seasonal Kendall



Figure E1110 Qu'Appelle River: Chromium Total

Time Series



Figure E1111 Qu'Appelle River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.76
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1112 Qu'Appelle River: Cobalt Dissolved



Figure E1113 Qu'Appelle River: Cobalt Dissolved



Figure E1114 Qu'Appelle River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 44.17
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1115 Qu'Appelle River: Cobalt Total

Seasonal Kendall



Figure E1116 Qu'Appelle River: Cobalt Total



Figure E1117 Qu'Appelle River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 24.3
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 24.3
Adjusted Kruskal-Wallis statistic (H') = 24.3



Figure E1118 Qu'Appelle River: Copper Dissolved



Figure E1119 Qu'Appelle River: Copper Dissolved



Figure E1120 Qu'Appelle River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.64
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1121 Qu'Appelle River: Copper Total

Seasonal Kendall



Figure E1122 Qu'Appelle River: Copper Total

Time Series



Figure E1123 Qu'Appelle River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.57
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1124 Qu'Appelle River: Iron Dissolved



Figure E1125 Qu'Appelle River: Iron Dissolved



Figure E1126 Qu'Appelle River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.22
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1127 Qu'Appelle River: Iron Total

Seasonal Kendall



Figure E1128 Qu'Appelle River: Iron Total

Time Series



Figure E1129 Qu'Appelle River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.528
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1130 Qu'Appelle River: Lead Dissolved



Figure E1131 Qu'Appelle River: Lead Dissolved



Figure E1132 Qu'Appelle River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.59
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 45.59
Adjusted Kruskal-Wallis statistic (H') = 45.59



Figure E1133 Qu'Appelle River: Lead Total

Seasonal Kendall



Figure E1134 Qu'Appelle River: Lead Total



Figure E1135 Qu'Appelle River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.157
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1136 Qu'Appelle River: Lithium Dissolved



Figure E1137 Qu'Appelle River: Lithium Dissolved



Figure E1138 Qu'Appelle River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.757
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.757
Adjusted Kruskal-Wallis statistic (H') = 1.757



Figure E1139 Qu'Appelle River: Lithium Total

Sen's Slope Estimator



Figure E1140 Qu'Appelle River: Lithium Total

Time Series



Figure E1141 Qu'Appelle River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 10.98
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1142 Qu'Appelle River: Manganese Dissolved



Figure E1143 Qu'Appelle River: Manganese Dissolved



Figure E1144 Qu'Appelle River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.07
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1145 Qu'Appelle River: Manganese Total

Seasonal Kendall



Figure E1146 Qu'Appelle River: Manganese Total



Figure E1147 Qu'Appelle River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.113
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1148 Qu'Appelle River: Molybdenum Dissolved



Figure E1149 Qu'Appelle River: Molybdenum Dissolved



Figure E1150 Qu'Appelle River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.385
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1151 Qu'Appelle River: Molybdenum Total

Sen's Slope Estimator



Figure E1152 Qu'Appelle River: Molybdenum Total



Figure E1153 Qu'Appelle River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.887
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1154 Qu'Appelle River: Nickel Dissolved



Figure E1155 Qu'Appelle River: Nickel Dissolved



Figure E1156 Qu'Appelle River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 40.23
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 40.23
Adjusted Kruskal-Wallis statistic (H') = 40.23



Figure E1157 Qu'Appelle River: Nickel Total

Seasonal Kendall



Figure E1158 Qu'Appelle River: Nickel Total

Time Series



Figure E1159 Qu'Appelle River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.2
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 2 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.2
Adjusted Kruskal-Wallis statistic (H') = 1.2



Figure E1160 Qu'Appelle River: Selenium Dissolved



Figure E1161 Qu'Appelle River: Selenium Dissolved



Figure E1162 Qu'Appelle River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.762
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.762
Adjusted Kruskal-Wallis statistic (H') = 1.762



Figure E1163 Qu'Appelle River: Selenium Total

Sen's Slope Estimator



Figure E1164 Qu'Appelle River: Selenium Total

Time Series



Figure E1165 Qu'Appelle River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 26.38
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1166 Qu'Appelle River: Silver Total



Figure E1167 Qu'Appelle River: Silver Total



Figure E1168 Qu'Appelle River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.61
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 14.61
Adjusted Kruskal-Wallis statistic (H') = 14.61



Figure E1169 Qu'Appelle River: Thallium Dissolved

Seasonal Kendall



Figure E1170 Qu'Appelle River: Thallium Dissolved

Time Series



Figure E1171 Qu'Appelle River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 40.54
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1172 Qu'Appelle River: Thallium Total



Figure E1173 Qu'Appelle River: Thallium Total



Figure E1174 Qu'Appelle River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.454
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1175 Qu'Appelle River: Uranium Dissolved

Sen's Slope Estimator



Figure E1176 Qu'Appelle River: Uranium Dissolved

Time Series



Figure E1177 Qu'Appelle River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.07585
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1178 Qu'Appelle River: Uranium Total



Figure E1179 Qu'Appelle River: Uranium Total



Figure E1180 Qu'Appelle River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.06
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1181 Qu'Appelle River: Vanadium Dissolved

Seasonal Kendall



Figure E1182 Qu'Appelle River: Vanadium Dissolved



Figure E1183 Qu'Appelle River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 55.93
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1184 Qu'Appelle River: Vanadium Total



Figure E1185 Qu'Appelle River: Vanadium Total



Figure E1186 Qu'Appelle River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 13.9
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 13.9
Adjusted Kruskal-Wallis statistic (H') = 13.9



Figure E1187 Qu'Appelle River: Zinc Dissolved

Seasonal Kendall



Figure E1188 Qu'Appelle River: Zinc Dissolved

Time Series



Figure E1189 Qu'Appelle River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 25.5
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 25.5
Adjusted Kruskal-Wallis statistic (H') = 25.5



Figure E1190 Qu'Appelle River: Zinc Total



Figure E1191 Qu'Appelle River: Zinc Total



Figure E1192 Red Deer River (SK-MB): Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.928
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1193 Red Deer River (SK-MB): Aluminum Dissolved



Figure E1194 Red Deer River (SK-MB): Aluminum Dissolved



Figure E1195 Red Deer River (SK-MB): Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.18
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1196 Red Deer River (SK-MB): Aluminum Total

Seasonal Kendall



Figure E1197 Red Deer River (SK-MB): Aluminum Total



Figure E1198 Red Deer River (SK-MB): Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 39.72
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1199 Red Deer River (SK-MB): Arsenic Dissolved



Figure E1200 Red Deer River (SK-MB): Arsenic Dissolved



Figure E1201 Red Deer River (SK-MB): Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.149
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1202 Red Deer River (SK-MB): Arsenic Total

Seasonal Kendall



Figure E1203 Red Deer River (SK-MB): Arsenic Total



Figure E1204 Red Deer River (SK-MB): Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.794
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1205 Red Deer River (SK-MB): Barium Dissolved



Figure E1206 Red Deer River (SK-MB): Barium Dissolved



Figure E1207 Red Deer River (SK-MB): Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 29.67
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1208 Red Deer River (SK-MB): Barium Total

Seasonal Kendall



Figure E1209 Red Deer River (SK-MB): Barium Total

Time Series



Figure E1210 Red Deer River (SK-MB): Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2098
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1211 Red Deer River (SK-MB): Beryllium Dissolved



Figure E1212 Red Deer River (SK-MB): Beryllium Dissolved



Figure E1213 Red Deer River (SK-MB): Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 22.89
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1214 Red Deer River (SK-MB): Beryllium Total

Seasonal Kendall



Figure E1215 Red Deer River (SK-MB): Beryllium Total



Figure E1216 Red Deer River (SK-MB): Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.01125
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1217 Red Deer River (SK-MB): Boron Dissolved



Figure E1218 Red Deer River (SK-MB): Boron Dissolved



Figure E1219 Red Deer River (SK-MB): Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1145
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1220 Red Deer River (SK-MB): Boron Total

Sen's Slope Estimator



Figure E1221 Red Deer River (SK-MB): Boron Total

Time Series



Figure E1222 Red Deer River (SK-MB): Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.683
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 3.683
Adjusted Kruskal-Wallis statistic (H') = 3.683



Figure E1223 Red Deer River (SK-MB): Cadmium Dissolved



Figure E1224 Red Deer River (SK-MB): Cadmium Dissolved



Figure E1225 Red Deer River (SK-MB): Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.07
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1226 Red Deer River (SK-MB): Cadmium Total

Seasonal Kendall



Figure E1227 Red Deer River (SK-MB): Cadmium Total



Figure E1228 Red Deer River (SK-MB): Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.344
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1229 Red Deer River (SK-MB): Chromium Dissolved



Figure E1230 Red Deer River (SK-MB): Chromium Dissolved



Figure E1231 Red Deer River (SK-MB): Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.23
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1232 Red Deer River (SK-MB): Chromium Total

Seasonal Kendall



Figure E1233 Red Deer River (SK-MB): Chromium Total



Figure E1234 Red Deer River (SK-MB): Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.191
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1235 Red Deer River (SK-MB): Cobalt Dissolved



Figure E1236 Red Deer River (SK-MB): Cobalt Dissolved



Figure E1237 Red Deer River (SK-MB): Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4201
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1238 Red Deer River (SK-MB): Cobalt Total

Sen's Slope Estimator



Figure E1239 Red Deer River (SK-MB): Cobalt Total

Time Series



Figure E1240 Red Deer River (SK-MB): Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.029
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1241 Red Deer River (SK-MB): Copper Dissolved



Figure E1242 Red Deer River (SK-MB): Copper Dissolved



Figure E1243 Red Deer River (SK-MB): Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.307
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1244 Red Deer River (SK-MB): Copper Total

Seasonal Kendall



Figure E1245 Red Deer River (SK-MB): Copper Total

Time Series



Figure E1246 Red Deer River (SK-MB): Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.9
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1247 Red Deer River (SK-MB): Iron Dissolved



Figure E1248 Red Deer River (SK-MB): Iron Dissolved



Figure E1249 Red Deer River (SK-MB): Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 36.63
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1250 Red Deer River (SK-MB): Iron Total

Seasonal Kendall



Figure E1251 Red Deer River (SK-MB): Iron Total

Time Series



Figure E1252 Red Deer River (SK-MB): Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.6254
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1253 Red Deer River (SK-MB): Lead Dissolved



Figure E1254 Red Deer River (SK-MB): Lead Dissolved



Figure E1255 Red Deer River (SK-MB): Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.404
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1256 Red Deer River (SK-MB): Lead Total

Seasonal Kendall



Figure E1257 Red Deer River (SK-MB): Lead Total

Time Series



Figure E1258 Red Deer River (SK-MB): Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.832
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1259 Red Deer River (SK-MB): Lithium Dissolved



Figure E1260 Red Deer River (SK-MB): Lithium Dissolved



Figure E1261 Red Deer River (SK-MB): Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.399
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1262 Red Deer River (SK-MB): Lithium Total

Sen's Slope Estimator



Figure E1263 Red Deer River (SK-MB): Lithium Total



Figure E1264 Red Deer River (SK-MB): Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.68
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1265 Red Deer River (SK-MB): Manganese Dissolved



Figure 1266 Red Deer River (SK-MB): Manganese Dissolved



Figure E1267 Red Deer River (SK-MB): Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4709
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1268 Red Deer River (SK-MB): Manganese Total

Sen's Slope Estimator



Figure E1269 Red Deer River (SK-MB): Manganese Total

Time Series



Figure E1270 Red Deer River (SK-MB): Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.697
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1271 Red Deer River (SK-MB): Molybdenum Dissolved



Figure E1272 Red Deer River (SK-MB): Molybdenum Dissolved



Figure E1273 Red Deer River (SK-MB): Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 6.379
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 6.379
Adjusted Kruskal-Wallis statistic (H') = 6.379



Figure E1274 Red Deer River (SK-MB): Molybdenum Total

Seasonal Kendall



Figure E1275 Red Deer River (SK-MB): Molybdenum Total

Time Series



Figure E1276 Red Deer River (SK-MB): Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.2011
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1277 Red Deer River (SK-MB): Nickel Dissolved



Figure E1278 Red Deer River (SK-MB): Nickel Dissolved



Figure E1279 Red Deer River (SK-MB): Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.985
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1280 Red Deer River (SK-MB): Nickel Total

Seasonal Kendall



Figure E1281 Red Deer River (SK-MB): Nickel Total



Figure E1282 Red Deer River (SK-MB): Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.166
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1283 Red Deer River (SK-MB): Selenium Dissolved



Figure E1284 Red Deer River (SK-MB): Selenium Dissolved



Figure E1285 Red Deer River (SK-MB): Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.04599
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1286 Red Deer River (SK-MB): Selenium Total

Sen's Slope Estimator



Figure E1287 Red Deer River (SK-MB): Selenium Total

Time Series



Figure E1288 Red Deer River (SK-MB): Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.764
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1289 Red Deer River (SK-MB): Silver Total



Figure E1290 Red Deer River (SK-MB): Silver Total



Figure E1291 Red Deer River (SK-MB): Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.56
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.56
Adjusted Kruskal-Wallis statistic (H') = 1.56



Figure E1292 Red Deer River (SK-MB): Thallium Dissolved

Sen's Slope Estimator



Figure E1293 Red Deer River (SK-MB): Thallium Dissolved

Time Series



Figure E1294 Red Deer River (SK-MB): Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.911
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1295 Red Deer River (SK-MB): Thallium Total



Figure E1296 Red Deer River (SK-MB): Thallium Total



Figure E1297 Red Deer River (SK-MB): Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.537
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1298 Red Deer River (SK-MB): Uranium Dissolved

Seasonal Kendall



Figure E1299 Red Deer River (SK-MB): Uranium Dissolved

Time Series



Figure E1300 Red Deer River (SK-MB): Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.046
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1301 Red Deer River (SK-MB): Uranium Total



Figure E1302 Red Deer River (SK-MB): Uranium Total



Figure E1303 Red Deer River (SK-MB): Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 34.83
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1304 Red Deer River (SK-MB): Vanadium Dissolved

Seasonal Kendall



Figure E1305 Red Deer River (SK-MB): Vanadium Dissolved



Figure E1306 Red Deer River (SK-MB): Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.03
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1307 Red Deer River (SK-MB): Vanadium Total



Figure E1308 Red Deer River (SK-MB): Vanadium Total



Figure E1309 Red Deer River (SK-MB): Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.25
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1310 Red Deer River (SK-MB): Zinc Dissolved

Seasonal Kendall



Figure E1311 Red Deer River (SK-MB): Zinc Dissolved

Time Series



Figure E1312 Red Deer River (SK-MB): Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 35.66
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1313 Red Deer River (SK-MB): Zinc Total



Figure E1314 Red Deer River (SK-MB): Zinc Total

Time Series



Figure E1315 Saskatchewan River: Aluminum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 28.08
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1316 Saskatchewan River: Aluminum Dissolved



Figure E1317 Saskatchewan River: Aluminum Dissolved



Figure E1318 Saskatchewan River: Aluminum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 46.14
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1319 Saskatchewan River: Aluminum Total

Seasonal Kendall



Figure E1320 Saskatchewan River: Aluminum Total

Time Series



Figure E1321 Saskatchewan River: Arsenic Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 16.31
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1322 Saskatchewan River: Arsenic Dissolved



Figure E1323 Saskatchewan River: Arsenic Dissolved



Figure E1324 Saskatchewan River: Arsenic Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.18
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1325 Saskatchewan River: Arsenic Total

Seasonal Kendall



Figure E1326 Saskatchewan River: Arsenic Total

Time Series



Figure E1327 Saskatchewan River: Barium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.88
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1328 Saskatchewan River: Barium Dissolved



Figure E1329 Saskatchewan River: Barium Dissolved



Figure E1330 Saskatchewan River: Barium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.5193
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1331 Saskatchewan River: Barium Total

Sen's Slope Estimator



Figure E1332 Saskatchewan River: Barium Total

Time Series



Figure E1333 Saskatchewan River: Beryllium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.4103
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1334 Saskatchewan River: Beryllium Dissolved



Figure E1335 Saskatchewan River: Beryllium Dissolved



Figure E1336 Saskatchewan River: Beryllium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 49.53
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1337 Saskatchewan River: Beryllium Total

Seasonal Kendall



Figure E1338 Saskatchewan River: Beryllium Total

Time Series



Figure E1339 Saskatchewan River: Boron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 5.648
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1340 Saskatchewan River: Boron Dissolved



Figure E1341 Saskatchewan River: Boron Dissolved



Figure E1342 Saskatchewan River: Boron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 4.105
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1343 Saskatchewan River: Boron Total

Seasonal Kendall



Figure E1344 Saskatchewan River: Boron Total

Time Series



Figure E1345 Saskatchewan River: Cadmium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 1.93
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 1.93
Adjusted Kruskal-Wallis statistic (H') = 1.93



Figure E1346 Saskatchewan River: Cadmium Dissolved



Figure E1347 Saskatchewan River: Cadmium Dissolved



Figure E1348 Saskatchewan River: Cadmium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 15.98
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1349 Saskatchewan River: Cadmium Total

Seasonal Kendall



Figure E1350 Saskatchewan River: Cadmium Total

Time Series



Figure E1351 Saskatchewan River: Chromium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 7.004
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1352 Saskatchewan River: Chromium Dissolved



Figure E1353 Saskatchewan River: Chromium Dissolved



Figure E1354 Saskatchewan River: Chromium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 32.86
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 1 groups of ties in the data, consequently the Kruskal-Wallis statistic (H) was adjusted. The adjusted statistic (H') was utilized to determine if the medians were equal.
Kruskal-Wallis statistic (H) = 32.86
Adjusted Kruskal-Wallis statistic (H') = 32.86



Figure E1355 Saskatchewan River: Chromium Total

Seasonal Kendall



Figure E1356 Saskatchewan River: Chromium Total



Figure E1357 Saskatchewan River: Cobalt Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 17.15
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1358 Saskatchewan River: Cobalt Dissolved



Figure E1359 Saskatchewan River: Cobalt Dissolved



Figure E1360 Saskatchewan River: Cobalt Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 44.14
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1361 Saskatchewan River: Cobalt Total

Seasonal Kendall



Figure E1362 Saskatchewan River: Cobalt Total

Time Series



Figure E1363 Saskatchewan River: Copper Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 19.72
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1364 Saskatchewan River: Copper Dissolved



Figure E1365 Saskatchewan River: Copper Dissolved



Figure E1366 Saskatchewan River: Copper Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 45.48
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1367 Saskatchewan River: Copper Total

Seasonal Kendall



Figure E1368 Saskatchewan River: Copper Total

Time Series



Figure E1369 Saskatchewan River: Iron Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 52.89
Tabulated Chi-Squared value = 7.815 with 3 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1370 Saskatchewan River: Iron Dissolved



Figure E1371 Saskatchewan River: Iron Dissolved



Figure E1372 Saskatchewan River: Iron Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 41.85
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1373 Saskatchewan River: Iron Total

Seasonal Kendall



Figure E1374 Saskatchewan River: Iron Total

Time Series



Figure E1375 Saskatchewan River: Lead Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 0.1241
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1376 Saskatchewan River: Lead Dissolved



Figure E1377 Saskatchewan River: Lead Dissolved



Figure E1378 Saskatchewan River: Lead Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 42.02
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1379 Saskatchewan River: Lead Total

Seasonal Kendall



Figure E1380 Saskatchewan River: Lead Total

Time Series



Figure E1381 Saskatchewan River: Lithium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 11.56
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1382 Saskatchewan River: Lithium Dissolved



Figure E1383 Saskatchewan River: Lithium Dissolved



Figure E1384 Saskatchewan River: Lithium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 3.885
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1385 Saskatchewan River: Lithium Total

Seasonal Kendall



Figure E1386 Saskatchewan River: Lithium Total



Figure E1387 Saskatchewan River: Manganese Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 70.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1388 Saskatchewan River: Manganese Dissolved



Figure E1389 Saskatchewan River: Manganese Dissolved



Figure E1390 Saskatchewan River: Manganese Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 9.124
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1391 Saskatchewan River: Manganese Total

Seasonal Kendall



Figure E1392 Saskatchewan River: Manganese Total

Time Series



Figure E1393 Saskatchewan River: Molybdenum Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 8.835
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1394 Saskatchewan River: Molybdenum Dissolved



Figure E1395 Saskatchewan River: Molybdenum Dissolved



Figure E1396 Saskatchewan River: Molybdenum Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 12.91
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1397 Saskatchewan River: Molybdenum Total

Seasonal Kendall



Figure E1398 Saskatchewan River: Molybdenum Total

Time Series



Figure E1399 Saskatchewan River: Nickel Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.88
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1400 Saskatchewan River: Nickel Dissolved



Figure E1401 Saskatchewan River: Nickel Dissolved



Figure E1402 Saskatchewan River: Nickel Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 33.98
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1403 Saskatchewan River: Nickel Total

Seasonal Kendall



Figure E1404 Saskatchewan River: Nickel Total

Time Series



Figure E1405 Saskatchewan River: Selenium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 23.43
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1406 Saskatchewan River: Selenium Dissolved



Figure E1407 Saskatchewan River: Selenium Dissolved



Figure E1408 Saskatchewan River: Selenium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 14.1
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1409 Saskatchewan River: Selenium Total

Seasonal Kendall



Figure E1410 Saskatchewan River: Selenium Total

Time Series



Figure E1411 Saskatchewan River: Silver Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 28.05
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1412 Saskatchewan River: Silver Total



Figure E1413 Saskatchewan River: Silver Total



Figure E1414 Saskatchewan River: Thallium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates NO SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is less than or equal to the Chi-squared value, we conclude that no season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 2.852
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1415 Saskatchewan River: Thallium Dissolved

Sen's Slope Estimator



Figure E1416 Saskatchewan River: Thallium Dissolved



Figure E1417 Saskatchewan River: Thallium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 40.52
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1418 Saskatchewan River: Thallium Total



Figure E1419 Saskatchewan River: Thallium Total



Figure E1420 Saskatchewan River: Uranium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 27.95
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1421 Saskatchewan River: Uranium Dissolved

Seasonal Kendall



Figure E1422 Saskatchewan River: Uranium Dissolved

Time Series



Figure E1423 Saskatchewan River: Uranium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 20.03
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1424 Saskatchewan River: Uranium Total



Figure E1425 Saskatchewan River: Uranium Total



Figure E1426 Saskatchewan River: Vanadium Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 38.87
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1427 Saskatchewan River: Vanadium Dissolved

Seasonal Kendall



Figure E1428 Saskatchewan River: Vanadium Dissolved

Time Series



Figure E1429 Saskatchewan River: Vanadium Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 54.33
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1430 Saskatchewan River: Vanadium Total



Figure E1431 Saskatchewan River: Vanadium Total



Figure E1432 Saskatchewan River: Zinc Dissolved

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 26.07
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1433 Saskatchewan River: Zinc Dissolved

Seasonal Kendall



Figure E1434 Saskatchewan River: Zinc Dissolved

Time Series



Figure E1435 Saskatchewan River: Zinc Total

Seasonality

For the selected data, the Kruskal-Wallis test indicates SEASONALITYat the 5% significance level. Because the calculated Kruskal-Wallis statistic is greater than the Chi-squared value, we conclude that at least one season has a significantly different median concentration of this constituent than any other season.
Calculated Kruskal-Wallis statistic = 35.12
Tabulated Chi-Squared value = 3.841 with 1 degrees of freedom at the 5% significance level.
There were 0 groups of ties in the data, so no adjustment to the Kruskal-Wallis statistic (H) was necessary.



Figure E1436 Saskatchewan River: Zinc Total



Figure E1437 Saskatchewan River: Zinc Total